

Ising model on nonorientable surfaces: Exact solution for the Möbius strip and the Klein bottle

Wentao T. Lu and F. Y. Wu

Department of Physics

Northeastern University, Boston, Massachusetts 02115

(February 6, 2008)

Closed-form expressions are obtained for the partition function of the Ising model on an $\mathcal{M} \times \mathcal{N}$ simple-quartic lattice embedded on a Möbius strip and a Klein bottle for finite $\mathcal{M}$ and $\mathcal{N}$. The finite-size effects at criticality are analyzed and compared with those under cylindrical and toroidal boundary conditions. Our analysis confirms that the central charge is $c = 1/2$.

I. INTRODUCTION

There has been considerable recent interest [1–3] in studying lattice models on nonorientable surfaces, both as new challenging unsolved lattice-statistical problems and as a realization and testing of predictions of the conformal field theory [4]. In a recent paper [1] we have presented the solution of dimers on the Möbius strip and Klein bottle and studied its finite-size corrections. In this paper we consider the Ising model.

The Ising model in two dimensions was first solved by Onsager in 1944 [5] who obtained the close-form expression of the partition function for a simple-quartic $\mathcal{M} \times \mathcal{N}$ lattice wrapped on a cylinder. The exact solution for an $\mathcal{M} \times \mathcal{N}$ lattice on a torus, namely, with periodic boundary condition in both directions, was obtained by Kaufman 4 years later [6]. Onsager and Kaufman used spinor analysis to derive the solutions, and the solution under the cylindrical boundary condition was rederived later by McCoy and Wu [7] using the method of dimers. As far as we know, these are the only known solutions of the two-dimensional Ising model on finite lattices. Here, using the method of dimers, we derive exact expressions for the partition function of the Ising model on finite Möbius strips and Klein bottles. As we shall see, as a consequence of the Möbius topology, the solution assumes a form which depends on whether the width of the lattice is even or odd. However, all solutions yield the same bulk free energy. We also present results of finite-size analyses for corrections to the bulk solution, and compare with those deduced under other boundary conditions. Our explicit calculations confirm that the central charge is $c = 1/2$.

II. THE $2M \times N$ MÖBIUS STRIP

To begin with, we consider a $2M \times N$ simple-quartic Ising lattice $\mathcal{L}$ embedded on a Möbius strip, where M, N are integers and $2M$ is the width of the strip. The ex-

ample of a lattice $\mathcal{L}$ for $2M = 4, N = 5$ is shown in Fig. 1.

While we shall consider the case of a uniform reduced interaction K , to facilitate considerations it is convenient to let the N vertical edges located in the middle of the strip to take on a different interaction K_1 as shown. By setting $K_1 = 0$ the Möbius strip reduces to an $M \times 2N$ strip with a “cylindrical” boundary condition, namely, periodic in one direction and free in the other, for which the partition function has been evaluated by McCoy and Wu [7]. By setting $K_1 = \infty$ the two center rows of spins coalesce into a single row with an (additive) interaction which in this case is $2K$. These are two key elements of our consideration.




FIG. 1. A 4×5 Möbius strip $\mathcal{L}$. Vertices labeled A, B, C, D are repeated sites.

Following standard procedures [7] we write the partition function of the Ising model on $\mathcal{L}$ as

$$Z_{2M,N}^{\text{Mob}}(K, K_1) = 2^{2MN} (\cosh K)^{2(2MN-N)} (\cosh K_1)^N \times G(z, z_1), \quad (1)$$

where $z = \tanh K$, $z_1 = \tanh K_1$, and

$$G(z, z_1) = \sum_{\text{closed polygons}} z^n z_1^{n_1} \quad (2)$$

is the generating function of all closed polygonal graphs on $\mathcal{L}$ with edge weights z and z_1 . Here, n is the number

of polygon edges with weight z and n_1 the number of edges with weight z_1 .

The generating function $G(z, z_1)$ is a multinomial in z and z_1 and, due to the Möbius topology, the integer n_1 can take on any value in $\{0, N\}$. Thus, we have

$$G(z, z_1) = \sum_{n_1=0}^N T_{n_1}(z) z_1^{n_1}, \quad (3)$$

where $T_{n_1}(z)$ are polynomials in z with strictly positive coefficients.

To evaluate $G(z, z_1)$, we again follow the usual procedure of mapping polygonal configurations on $\mathcal{L}$ onto dimer configurations on a dimer lattice $\mathcal{L}_D$ of $8MN$ sites, constructed by expanding each site of $\mathcal{L}$ into a “city” of 4 sites [7,8]. The resulting $\mathcal{L}_D$ for the 4×5 $\mathcal{L}$ is shown in Fig. 2.




FIG. 2. The dimer lattice $\mathcal{L}_D$ corresponding to the 4×5 Möbius strip.

As the deletion of all z_1 edges reduces the lattice to one with a cylindrical boundary condition solved in [7], we orient all edges of weights z and 1 as in [7]. In addition, all z_1 edges are oriented in the direction as shown in Fig. 2. Then, we have the following:

Theorem:

Let A be the $8MN \times 8MN$ antisymmetric determinant defined by the lattice edge orientation shown in Fig. 2, and let

$$\text{Pf}A(z, z_1) = \sqrt{\det A(z, z_1)} \quad (4)$$

denote the Pfaffian of A . Then

$$\text{Pf}A(z, z_1) = \sum_{n_1=0}^N \epsilon_{n_1} T_{n_1}(z) z_1^{n_1}, \quad (5)$$

where $\epsilon_{4m} = \epsilon_{4m+1} = 1$, $\epsilon_{4m+2} = \epsilon_{4m+3} = -1$ for any integer $m \geq 0$.

Remark: Define

$$X_p = \sum_{m=0}^{[N/4]} T_{4m+p}(z) z_1^{4m+p}, \quad p = 0, 1, 2, 3, \quad (6)$$

where $[N/4]$ is the integral part of $N/4$ so that $G(z, z_1) = X_0 + X_1 + X_2 + X_3$. It then follows from (5) that we have

$$\text{Pf}A(z, \pm iz_1) = X_0 + X_2 \pm i(X_1 + X_3). \quad (7)$$

As a consequence, we obtain

$$G(z, z_1) = \frac{1}{2} \left[(1 - i) \text{Pf}A(z, iz_1) + (1 + i) \text{Pf}A(z, -iz_1) \right], \quad (8)$$

where, as evaluated in the next section, the Pfaffian is given by

$$\text{Pf}A(z, z_1) = [z(1 - z^2)]^{MN} \times \prod_{n=1}^N \left[\frac{\sinh(M+1)t(\phi_n) - c(z, z_1) \sinh Mt(\phi_n)}{\sinh t(\phi_n)} \right], \quad (9)$$

with

$$\begin{aligned} c(z, z_1) &= \frac{z(1 + z^2 + 2z \cos \phi_n) + 2(-1)^n z_1 \sin \phi_n}{1 - z^2} \\ \cosh t(\phi) &= \cosh 2K \coth 2K - \cos \phi \\ \phi_n &= (2n - 1)\pi/2N. \end{aligned} \quad (10)$$

Here we have used the fact that $\prod_{n=1}^N = \prod_{n=N+1}^{2N}$ in the product in (9). Substituting these results into (1), we are led to the following explicit expression for the partition function,

$$Z_{2M,N}^{\text{Mob}}(K, K) = \frac{1}{2} \left(2 \sinh 2K \right)^{MN} \times \left[(1 - i)F_+ + (1 + i)F_- \right], \quad (11)$$

where

$$F_{\pm} = \prod_{n=1}^N \left[e^{Mt(\phi_n)} \left(\frac{e^{t(\phi_n)} - c(z, \pm iz_1)}{2 \sinh t(\phi_n)} \right) - e^{-Mt(\phi_n)} \left(\frac{e^{-t(\phi_n)} - c(z, \pm iz_1)}{2 \sinh t(\phi_n)} \right) \right]. \quad (12)$$

This completes the evaluation of the Ising partition function for the $2M \times N$ Möbius strip. Note that we have $\cosh t(\phi_n) \geq 1$ so we can always take $t(\phi_n) \geq 0$. The leading contribution in (12) for large M is therefore $\prod e^{Mt(\phi_n)}$.

For the 2×5 Möbius strip, for example, we find

$$\begin{aligned} \text{Pf}A(z, z_1) &= 1 + z^{10} + 10z_1 z^5 - 5z_1^2 z^2 (1 + z^2 + z^4 + z^6) \\ &\quad - 20z_1^3 z^5 + 5z_1^4 z^4 (1 + z^2) + 2z_1^5 z^5, \\ G(z, z_1) &= 1 + z^{10} + 10z_1 z^5 + 5z_1^2 z^2 (1 + z^2 + z^4 + z^6) \\ &\quad + 20z_1^3 z^5 + 5z_1^4 z^4 (1 + z^2) + 2z_1^5 z^5, \end{aligned} \quad (13)$$

which can be verified by explicit enumerations.

We next prove the theorem.

Considered as a multinomial in z and z_1 , there exists a one-one correspondence between terms in the dimer generating function $G(z, z_1)$ and (combinations of) terms in the Pfaffian (4). However, while all terms in $G(z, z_1)$ are positive, terms in the Pfaffian do not necessarily possess the same sign. The crux of the matter is to find an appropriate linear combination of Pfaffians to yield the desired $G(z, z_1)$. For this purpose it is convenient to compare an arbitrary term C_1 in the Pfaffian with a standard one C_0 . We choose C_0 to be one in which no z and z_1 dimers are present.

The superposition of two dimer configurations represented by C_0 and C_1 produces superposition polygons. Kasteleyn [9] has shown that the two terms will have the same sign if all superposition polygons are oriented “clockwise-odd”, namely, there is an odd number of edges oriented in the clockwise direction.

Now since all z and 1 edges of $\mathcal{L}_D$ are oriented as in [7], terms in the Pfaffian with no z_1 edges ($n_1 = 0$) will have the same sign as C_0 . To determine the sign of a term when z_1 edges are present, we associate a $+$ sign to each clockwise-odd superposition polygon and a $-$ sign to each clockwise-even superposition polygon. Then the sign of C_1 relative to C_0 is the product of the signs of all superposition polygons. The following elementary facts can be readily verified:

- i) Deformations of the borders of a superposition polygon always change m_1 , the number of its z_1 edges, by multiples of 2.
- ii) The sign of a superposition polygon is reversed under border deformations which change m_1 by 2.
- iii) Superposition polygons having 0 or 1 z_1 edges have a sign $+$.
- iv) There can be at most one superposition polygon having an odd number of z_1 edges (a property unique to nonorientable surfaces).

Let $m_1 = 4m + p$, where m is an integer and $p = 0, 1, 2, 3$. Because of iv), we need only to consider the presence of at most one polygon having $p = 1$ or 3. It now follows from i) and iii) that $\epsilon_{4m} = \epsilon_{4m+1} = +$, and from i), ii) and iii) that $\epsilon_{4m+2} = \epsilon_{4m+3} = -$. This establishes the theorem.

III. EVALUATION OF THE PFAFFIAN

In this section we derive the expression (9).

From the edge orientation of $\mathcal{L}_D$ of Fig. 2, one finds that the $8MN \times 8MN$ antisymmetric matrix A assumes the form

$$A(z, z_1) = A_0(z) \otimes I_{2N} + A_+(z) \otimes J_{2N} + A_-(z) \otimes J_{2N}^T + A_1(z_1) \otimes H_{2N}, \quad (14)$$

where A_0, A_+, A_-, A_1 are $4M \times 4M$ matrices, I_{2N} is the $2N \times 2N$ identity matrix, and J_{2N}, H_{2N} are the $2N \times 2N$ matrices

$$J_{2N} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -1 & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad H_{2N} = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}. \quad (15)$$

In addition, one has

$$\begin{aligned} A_0(z) &= a_{0,0} \otimes I_M + a_{0,1}(z) \otimes F_M + a_{0,-1}(z) \otimes F_M^T \\ A_\pm(z) &= a_{\pm 1,0}(z) \otimes I_M \\ A_1(z_1) &= a(z_1) \otimes G_M, \end{aligned} \quad (16)$$

where F_M, G_M are $M \times M$ matrices

$$\begin{aligned} F_M &= \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \\ G_M &= \begin{pmatrix} 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix}, \end{aligned} \quad (17)$$

F_M^T is the transpose of F_M , and

$$\begin{aligned} a_{0,0} &= \begin{pmatrix} 0 & 1 & -1 & -1 \\ -1 & 0 & 1 & -1 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & 0 \end{pmatrix}, \\ a(z_1) &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & z_1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ a_{1,0}(z) &= \begin{pmatrix} 0 & z & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ a_{0,1}(z) &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & z \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ a_{-1,0}(z) &= a_{1,0}^T(z) \\ a_{0,-1}(z) &= a_{0,1}^T(z). \end{aligned} \quad (18)$$

We use the fact that the determinant in (4) is equal to the product of the eigenvalues of the matrix A . To evaluate the latter, we note that J_{2N}, J_{2N}^T and H_{2N} mutually commute so that they can be diagonalized simultaneously. This leads to the respective eigenvalues $e^{i\phi_n}, e^{-i\phi_n}$ and $i(-1)^{n+1}$ and the expression

$$\det A(z, z_1) = \prod_{n=1}^{2N} \det A_M(z, z_1; \phi_n), \quad (19)$$

where

$$A_M(z, z_1; \phi_n) = A_0(z) + A_+(z)e^{i\phi_n} + A_-(z)e^{-i\phi_n} + i(-1)^{n+1}A_1(z_1) \quad (20)$$

is a $4M \times 4M$ matrix. Writing out explicitly, we have

$$A_M(z, z_1; \phi_n) = \begin{pmatrix} B(z) & a_{0,1}(z) & 0 & 0 \\ a_{0,-1}(z) & B(z) & a_{0,1}(z) & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & a_{0,-1}(z) & B(z) & a_{0,1}(z) \\ 0 & a_{0,-1}(z) & C(z, z_1) & 0 \end{pmatrix} \quad (21)$$

where $C(z, z_1) = B(z) + i(-1)^{n+1}a(z_1)$, and

$$B(z) = \begin{pmatrix} 0 & 1 + ze^{i\phi_n} & -1 & -1 \\ -(1 + ze^{-i\phi_n}) & 0 & 1 & -1 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & 0 \end{pmatrix}. \quad (22)$$

The evaluation of $\det A_M(z, z_1; \phi_n)$ can be carried out by using a recursion procedures introduced in [10] for a self-dual Ising model. Specifically, let $B_M = \det A_M(z, z_1; \phi_n)$ and D_M the determinant of the matrix $A_M(z, z_1; \phi_n)$ with the fourth row and fourth column removed. Then by expanding the determinants one finds the recursion relation (which is the same as that in [10])

$$\begin{pmatrix} B_M \\ D_M \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} B_{M-1} \\ D_{M-1} \end{pmatrix}, \quad M \geq 2, \quad (23)$$

with

$$\begin{aligned} a_{11} &= 1 + z^2 - 2z \cos \phi_n, \\ a_{12} &= -2iz^3 \sin \phi_n, \\ a_{21} &= 2iz \sin \phi_n, \\ a_{22} &= z^2(1 + z^2 + 2z \cos \phi_n), \end{aligned} \quad (24)$$

and the initial condition (which is different from [10])

$$\begin{aligned} B_1 &= B_1(z, z_1) \\ &\equiv 1 - 2z \cos \phi_n + z^2 - 2(-1)^n z z_1 \sin \phi_n, \\ D_1 &= D_1(z, z_1) \\ &\equiv 2iz \sin \phi_n - i(-1)^n z_1(1 + 2z \cos \phi_n + z^2). \end{aligned} \quad (25)$$

This leads to the solution

$$\begin{aligned} B_M &= B_1 \frac{\lambda_+^M - \lambda_-^M}{\lambda_+ - \lambda_-} - (a_{22}B_1 - a_{12}D_1) \frac{\lambda_+^{M-1} - \lambda_-^{M-1}}{\lambda_+ - \lambda_-}, \\ D_M &= D_1 \frac{\lambda_+^M - \lambda_-^M}{\lambda_+ - \lambda_-} - (a_{11}D_1 - a_{21}B_1) \frac{\lambda_+^{M-1} - \lambda_-^{M-1}}{\lambda_+ - \lambda_-}, \end{aligned} \quad (26)$$

where $\lambda_{\pm} = z(1 - z^2)e^{\pm t(\phi_n)}$ are the eigenvalues of the 2×2 matrix in (23). After some algebraic manipulation, this yields the expression (9) quoted in the preceding section.

IV. THE $(2M - 1) \times N$ MÖBIUS STRIP

We consider a $(2M - 1) \times N$ Möbius strip in this section.

In order to make use of results of the preceding sections, we start from the $2M \times N$ strip of Sec. 2, and let spins in the two center rows of the strip (the M th and $(M + 1)$ th rows) having interactions $K_0 = K/2$. The example of a 4×5 lattice with these interactions is shown in Fig. 3. Then, by taking $K_1 = \infty$ ($z_1 = 1$) as described in Sec. 1, this lattice reduces to the desired $(2M - 1) \times N$ Möbius strip of a uniform interaction K .




FIG. 3. Labelings of a 4×5 Möbius strip which reduces to a 3×5 Möbius strip upon taking $K_1 = \infty$.

Following this procedure, we have

$$Z_{2M-1,N}^{\text{Mob}}(K) = 2^{(2M-1)N} (\cosh K)^{4(M-1)N} \times \cosh^{2N}(K/2) G(z, z_0, z_1) \Big|_{z_1=1}, \quad (27)$$

where $z_0 = \tanh(K/2)$, and $G(z, z_0, z_1)$ is the generating function of closed polygons on the $2M \times N$ Möbius net with edge weights as described in the above.

The generating function $G(z, z_0, z_1)$ can be evaluated as in the previous sections. In place of (4), (19) and (21), we now have

$$\begin{aligned} \text{Pf} A(z, z_0, z_1) &= \sqrt{\det A(z, z_0, z_1)} \\ &= \prod_{n=1}^{2N} \sqrt{\det A_M(z, z_0, z_1; \phi_n)} \end{aligned} \quad (28)$$

with

$$A_M(z, z_0, z_1; \phi_n) = \begin{pmatrix} B(z) & a_{0,1}(z) & 0 & 0 \\ a_{0,-1}(z) & B(z) & a_{0,1}(z) & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & a_{0,-1}(z) & B(z) & a_{0,1}(z) \\ 0 & a_{0,-1}(z) & C(z, z_1) & 0 \end{pmatrix}. \quad (29)$$

Then (8) becomes

$$G(z, z_0, z_1) = \frac{1}{2} \left[(1-i) \text{Pf}A(z, z_0, iz_1) + (1+i) \text{Pf}A(z, z_0, -iz_1) \right]. \quad (30)$$

The evaluation of $\det A_M(z, z_0, z_1; \phi_n)$ can again be done recursively. Define as before $B_M = \det A_M(z, z_0, z_1; \phi_n)$ and D_M the determinant of A_M with the fourth row and column removed, one obtains again the recursion relations (23) and arrives at precisely the same solution (26), but now with a different initial condition

$$B_1 = B_1(z_0, z_1), \quad D_1 = D_1(z_0, z_1), \quad (31)$$

where the functions B_1 and D_1 are defined in (25). After some algebra, this leads to

$$\text{Pf}A(z, z_0, z_1) = [z(1-z^2)]^{(M-1)N} \times \prod_{n=1}^N \left[\frac{c_1 \sinh Mt(\phi_n) - c_2 \sinh(M-1)t(\phi_n)}{\sinh t(\phi_n)} \right], \quad (32)$$

where

$$c_1 = \frac{2z_0}{z} \left\{ 1 - z \left[\cos \phi_n + (-1)^n z_1 \sin \phi_n \right] \right\}, \\ c_2 = 2z_0 \left\{ 1 + z \left[\cos \phi_n + (-1)^n z_1 \sin \phi_n \right] \right\}. \quad (33)$$

The substitution of (32) into (30) and (27) now completes the evaluation of the partition function for a $(2M-1) \times N$ Möbius strip.

V. THE KLEIN BOTTLE

The Ising model on a Klein bottle can be considered similarly. We consider first a $2M \times N$ lattice $\mathcal{L}$, constructed by connecting the upper and lower edges of the Möbius strip of Fig. 1 in a periodic fashion with N extra vertical edges. As in the case of the Möbius strip, it is convenient to let the extra edges have interactions K_2 . The solution for a uniform interaction K is obtained at the end by setting $K_1 = K_2 = K$.

The Ising partition function for the Klein bottle now assumes the form

$$Z_{2M,N}^{\text{Kln}}(K, K_1, K_2) = 2^{2MN} (\cosh K)^{4MN-2N} \times (\cosh K_1 \cosh K_2)^N G^{\text{Kln}}(z, z_1, z_2), \quad (34)$$

where

$$G^{\text{Kln}}(z, z_1, z_2) = \sum_{\text{closed polygons}} z^n z_1^{n_1} z_2^{n_2} \quad (35)$$

generates all closed polygons on the $2M \times N$ lattice $\mathcal{L}$ with edge weights $z = \tanh K$, $z_1 = \tanh K_1$, and $z_2 = \tanh K_2$. The desired partition function is given by

$$Z_{2M,N}^{\text{Kln}}(K, K, K) = 2^{2MN} (\cosh K)^{4MN} G^{\text{Kln}}(z, z, z). \quad (36)$$

Again, it is necessary to first write $G^{\text{Kln}}(z, z_1, z_2)$ as a multinomial in z, z_1, z_2 in the form of

$$G^{\text{Kln}}(z, z_1, z_2) = \sum_{m,n=0}^N T_{m,n}(z) z_1^m z_2^n, \quad (37)$$

where $T_{m,n}(z)$ are polynomials in z with strictly positive coefficients.

The evaluation of $G^{\text{Kln}}(z, z_1, z_2)$ parallels that of $G(z, z_1)$ for the Möbius strip. One first maps the lattice $\mathcal{L}$ into a dimer lattice $\mathcal{L}_D$ by expanding each site into a city of 4 sites as shown in Fig. 2. Orient all K and K_1 edges of $\mathcal{L}_D$ as shown, and orient all K_2 edges in the same (downward) direction as the K_1 edges. Then this defines an $8MN \times 8MN$ antisymmetric matrix obtained by adding an extra term to $A(z, z_1)$ given by (14), namely,

$$A^{\text{Kln}}(z, z_1, z_2) = A(z, z_1) + b(z_2) \otimes G'_M \otimes H_{2N}. \quad (38)$$

Here,

$$b(z_2) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -z_2 \end{pmatrix}, \\ G'_M = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \end{pmatrix}. \quad (39)$$

Then, in place of theorem (5), we now have

$$\text{Pf}A^{\text{Kln}}(z, z_1, z_2) = \sum_{m,n=0}^N \epsilon_m \epsilon_n T_{m,n}(z) z_1^m z_2^n, \quad (40)$$

from which one obtains in a similar manner the result

$$G^{\text{Kln}}(z, z_1, z_2) = \frac{1}{2} \left[\text{Pf}A^{\text{Kln}}(z, iz_1, -iz_2) + \text{Pf}A^{\text{Kln}}(z, -iz_1, iz_2) - i \text{Pf}A^{\text{Kln}}(z, iz_1, iz_2) + i \text{Pf}A^{\text{Kln}}(z, -iz_1, -iz_2) \right]. \quad (41)$$

To evaluate the Pfaffian (40), we note that the matrix (38) can again be diagonalized in the $\{2N\}$ subspace, yielding

$$\begin{aligned} \text{Pf}A^{\text{Kln}}(z, z_1, z_2) &= \sqrt{\det A^{\text{Kln}}(z, z_1, z_2)} \\ &= \prod_{n=1}^{2N} \sqrt{\det A_M^{\text{Kln}}(z, z_1, z_2; \phi_n)}, \end{aligned} \quad (42)$$

where

$$\begin{aligned} A_M^{\text{Kln}}(z, z_1, z_2; \phi_n) &= A_M(z, z_1; \phi_n) \\ &\quad + i(-1)^{n+1} b(z_2) \otimes G'_M. \end{aligned} \quad (43)$$

Now we expand $\det A_M^{\text{Kln}}$ in z_2 . Since setting $z_2 = 0$ the determinant is precisely B_M and the term linear in z_2 , the $\{4, 4\}$ element of the determinant, is by definition D_M , one obtains

$$\det A_M^{\text{Kln}}(z, z_1, z_2; \phi_n) = B_M + i(-1)^n z_2 D_M, \quad M \geq 2, \quad (44)$$

where B_M and D_M have been computed in (26). This leads to

$$\begin{aligned} \text{Pf}A^{\text{Kln}}(z, z_1, z_2) &= \left(1 + \frac{z_1 z_2}{z^2}\right)^N \left[z(1 - z^2)\right]^{MN} \\ &\quad \times \prod_{n=1}^N \left[\frac{\sinh(M+1)t - c(z, z_1, z_2) \sinh Mt}{\sinh t} \right] \end{aligned} \quad (45)$$

where

$$\begin{aligned} c(z, z_1, z_2) &= \frac{1}{z(1 - z^2)(z^2 + z_1 z_2)} \left[(1 + z^2)(z^4 + z_1 z_2) \right. \\ &\quad \left. + 2z(z^4 - z_1 z_2) \cos \phi_n \right. \\ &\quad \left. + 2(-1)^n (z_1 + z_2) z^3 \sin \phi_n \right]. \end{aligned} \quad (46)$$

Setting $z_1 = z_2 = z$ in (41) and using (45), we obtain after some algebra

$$\begin{aligned} G^{\text{Kln}}(z, z, z) &= \left[z(1 - z^2)\right]^{MN} \left[\prod_{n=1}^N 2 \cosh Mt(\phi_n) \right. \\ &\quad \left. + \text{Im} \prod_{n=1}^N \left(\frac{\sinh Mt(\phi_n)}{\sinh t(\phi_n)} D(\phi_n) \right) \right], \end{aligned} \quad (47)$$

where

$$\begin{aligned} D(\phi_n) &= \frac{1}{z(1 - z^2)} \left[(1 - z^4) - 2z(1 + z^2) \cos \phi_n \right. \\ &\quad \left. - 4i(-1)^n z^2 \sin \phi_n \right] \end{aligned} \quad (48)$$

and Im denotes the imaginary part. The substitution of (47) into (36) now completes the evaluation of the partition function for a $2M \times N$ Klein bottle.

For the 2×2 Klein bottle, for example, we find

$$\begin{aligned} \text{Pf}A^{\text{Kln}}(z, z_1, z_2) &= 1 + z^4 + 4(z_1 + z_2)z^2 - 2(z_1^2 + z_2^2)z^2 \\ &\quad + 2z_1 z_2 (1 + z^2)^2 - 4z_1 z_2 (z_1 + z_2)z^2 \\ &\quad + z_1^2 z_2^2 (1 + z^4), \\ G^{\text{Kln}}(z, z_1, z_2) &= 1 + z^4 + 4(z_1 + z_2)z^2 + 2(z_1^2 + z_2^2)z^2 \\ &\quad + 2z_1 z_2 (1 + z^2)^2 + 4z_1 z_2 (z_1 + z_2)z^2 \\ &\quad + z_1^2 z_2^2 (1 + z^4), \end{aligned} \quad (49)$$

which can be verified by explicit enumerations.

For a $(2M - 1) \times N$ Klein bottle we can proceed as before by first considering a $2M \times N$ Klein bottle with interactions K, K_1, K_2 and, within the center two rows, interactions $K_0 = K/2$ as shown in Fig. 3. This is followed by taking $K_1 \rightarrow \infty$ and $K_2 = K$. Thus, in place of (34), we have

$$\begin{aligned} Z_{2M-1, N}^{\text{Kln}}(K) &= 2^{(2M-1)N} (\cosh K)^{(4M-3)N} \\ &\quad \times \cosh^{2N}(K/2) G^{\text{Kln}}(z, z_0, 1, z), \end{aligned} \quad (50)$$

where $z_0 = \tanh(K/2)$ and $G(z, z_0, z_1, z_2)$ generates polygonal configurations on the $2M \times N$ lattice. Then, as in the above, we find

$$\begin{aligned} G^{\text{Kln}}(z, z_0, z_1, z_2) &= \\ \frac{1}{2} &\left[\text{Pf}A^{\text{Kln}}(z, z_0, iz_1, -iz_2) + \text{Pf}A^{\text{Kln}}(z, z_0, -iz_1, iz_2) \right. \\ &\quad \left. - i \text{Pf}A^{\text{Kln}}(z, z_0, iz_1, iz_2) + i \text{Pf}A^{\text{Kln}}(z, z_0, -iz_1, -iz_2) \right], \end{aligned} \quad (51)$$

where $\text{Pf}A^{\text{Kln}}(z, z_0, z_1, z_2)$ is found to be given by the right-hand side of (32), but with

$$\begin{aligned} c_1 &= (1 + z_0^2)(1 - z_1 z_2) - 2z_0(1 + z_1 z_2) \cos \phi_n \\ &\quad - 2(-1)^n (z_1 + z_2) z_0 \sin \phi_n, \\ c_2 &= \frac{1}{z(1 - z^2)} \left\{ (z^2 + z_1 z_2) \left[(1 - z_0 z)^2 + (z - z_0)^2 \right] \right. \\ &\quad \left. + 2(z - z_0)(1 - z_0 z) \left[(z^2 - z_1 z_2) \cos \phi_n \right. \right. \\ &\quad \left. \left. + (-1)^n (z^2 z_1 + z_2) \sin \phi_n \right] \right\}, \end{aligned} \quad (52)$$

expressions which are valid for arbitrary z, z_0, z_1 , and z_2 . For $z_0 = \tanh(K/2)$, the case we are considering, (52) reduces to

$$\begin{aligned} c_1 &= \frac{2z_0}{z} \left[1 - z_1 z_2 - z(1 + z_1 z_2) \cos \phi_n \right. \\ &\quad \left. - (-1)^n z(z_1 + z_2) \sin \phi_n \right], \\ c_2 &= \frac{2z_0}{z^2} \left[z^2 + z_1 z_2 + z(z^2 - z_1 z_2) \cos \phi_n \right. \\ &\quad \left. + (-1)^n z(z^2 z_1 + z_2) \sin \phi_n \right], \end{aligned} \quad (53)$$

which reduces further to (33) after setting $z_2 = 0$. The explicit expression for the partition function is now obtained by substituting (51) into (50).

VI. THE BULK LIMIT AND FINITE-SIZE CORRECTIONS

In the thermodynamic limit, our solutions of the Ising partition function give rise to a bulk free energy

$$f_{\text{bulk}}(K) = \lim_{M,N \rightarrow \infty} \frac{1}{2MN} \ln Z(K) \quad (54)$$

identical to that of the Onsager solution [5]. Here, $Z(K)$ is any one of the 4 partition functions. Indeed, using the solution (11) for the $2M \times N$ Möbius strip, for example, one obtains

$$\begin{aligned} f_{\text{bulk}}(K) &= C(K) + \lim_{N \rightarrow \infty} (2N)^{-1} \sum_{n=1}^N t(\phi_n) \\ &= C(K) + \frac{1}{2\pi} \int_0^\pi d\phi \, t(\phi) \\ &= C(K) + \frac{1}{2\pi^2} \int_0^\pi d\phi \int_0^\pi d\theta \\ &\quad \ln \left[2 \cosh 2K \coth 2K - 2(\cos \theta + \cos \phi) \right], \end{aligned} \quad (55)$$

where $C(K) = [\ln(2 \sinh 2K)]/2$. This expression is the Onsager solution (steps leading to the last line of (55) can be found in [11]). It is well-known that $f_{\text{bulk}}(K)$ is singular at the critical point $\sinh 2K_c = 1$ or $2K_c = \ln(\sqrt{2} + 1)$.

For large M and N , one can use the Euler-MacLaurin summation formula to evaluate corrections to the bulk free energy, an analysis first carried out by Ferdinand and Fisher [12] for the Kaufman solution of the Ising model on a torus. Generally, for large M and N , we expect to have

$$\begin{aligned} \ln Z_{2M,N}(K) &= 2MN f_{\text{bulk}}(K) + N c_1(\xi, K) \\ &\quad + 2M c_2(\xi, K) + c_3(\xi, K) + \dots \end{aligned} \quad (56)$$

where $\xi = N/2M$ is the aspect ratio of the lattice. For the purpose of comparing with the conformal field theory [4], it is of particular interest to analyze corrections at the critical point. Following [12] as well as similar analyses for dimer systems [1,13], we have carried out such analyses for our solutions as well as for the solution of the Ising model under cylindrical boundary conditions [5,7].

For the $2M \times N$ Möbius strip, for example, one starts with the explicit expression (11) of the partition function, and uses the Euler-MacLaurin formula to evaluate corrections to the bulk free energy. The analysis is lengthy, even at the critical point K_c . We shall give details elsewhere and quote here only the results:

$$\begin{aligned} c_1(\xi, K_c) &\equiv c_1^{\text{Mob}} = I - K_c = -0.087 \, 618 \, \dots, \\ c_2(\xi, K_c) &= 0, \\ c_3(\xi, K_c) &= -\frac{1}{2} \ln 2 + \frac{1}{12} \ln \left[\frac{2\vartheta_3^2(0|i\xi)}{\vartheta_2(0|i\xi)\vartheta_4(0|i\xi)} \right] \\ &\quad + \frac{1}{2} \ln \left[1 + \frac{\vartheta_3(0|i\xi/2) - \vartheta_4(0|i\xi/2)}{2\vartheta_3(0|i\xi)} \right] \end{aligned} \quad (57)$$

where

$$\begin{aligned} I &= \frac{1}{2\pi} \int_0^\pi \ln \left(\sqrt{2} \sin \phi + \sqrt{1 + \sin^2 \phi} \right) d\phi \\ &= 0.353 \, 068 \, \dots \end{aligned}$$

and $\vartheta_i(u|\tau)$, $i = 2, 3, 4$, are the Jacobi theta functions [14]

$$\begin{aligned} \vartheta_2(u|\tau) &= 2 \sum_{n=1}^{\infty} q^{(n-\frac{1}{2})^2} \cos(2n-1)u, \\ \vartheta_3(u|\tau) &= 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \cos 2nu, \\ \vartheta_4(u|\tau) &= 1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2} \cos 2nu, \end{aligned} \quad (58)$$

with $q = e^{i\pi\tau}$. For the $2M \times N$ Klein bottle, we find

$$\begin{aligned} c_1(\xi, K_c) &= 0 \\ c_2(\xi, K_c) &= 0 \\ c_3(\xi, K_c) &= \frac{1}{6} \ln \left[\frac{2\vartheta_3^2(0|2i\xi)}{\vartheta_2(0|2i\xi)\vartheta_4(0|2i\xi)} \right] \\ &\quad + \ln \left[1 + \sqrt{\frac{\vartheta_2(0|2i\xi)}{2\vartheta_3(0|2i\xi)}} \right]. \end{aligned} \quad (59)$$

If one takes the limit of $N \rightarrow \infty$ ($M \rightarrow \infty$) first while keeping M (N) finite, one obtains

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z_{2M,N}(K_c) &= 2M f_{\text{bulk}}(K_c) + c_1 \\ &\quad + \Delta_1/2M + O(1/M^2), \\ \lim_{M \rightarrow \infty} \frac{1}{2M} \ln Z_{2M,N}(K_c) &= N f_{\text{bulk}}(K_c) + c_2 \\ &\quad + \Delta_2/N + O(1/N^2), \end{aligned} \quad (60)$$

where $c_1, c_2, \Delta_1, \Delta_2$ are constants. We list our results in the table below. Also listed are values for the Ising model with toroidal boundary conditions taken from [12], and values for the cylindrical boundary conditions computed using the solution of [7]. Our values of Δ_1 and Δ_2 imply a central charge

$$c = 1/2 \quad (61)$$

for the Ising model. This is consistent with the conformal field theory prediction [4].

	Möbius	Klein	Cylindrical	Toroidal
c_1	c_1^{Mob}	0	c_1^{Mob}	0
c_2	0	0	0	0
Δ_1	$\pi/48$	$\pi/12$	$\pi/48$	$\pi/12$
Δ_2	$\pi/48$	$\pi/48$	$\pi/12$	$\pi/12$

VII. SUMMARY

We have solved and obtained close-form expressions for the partition function of an Ising model on finite Möbius strips and Klein bottles with a uniform interaction K . The solution assumes different forms depending on whether the width of the lattice is even or odd. For a $2M \times N$ Möbius strip, where $2M$ is its width, the partition function $Z_{2M,N}^{\text{Mob}}(K)$ is given by (11) with $F_{\pm}$ given by (12). For a $(2M-1) \times N$ Möbius strip we employ a trick by first considering a $2M \times N$ lattice and then “fusing” it into the desired lattice by coalescing two rows of spins. The resulting partition function $Z_{2M-1,N}^{\text{Mob}}(K)$ is given by (27) in which the generating function $G(z, z_0, z_1)$ is (30) with the Pfaffians given by (32).

For a $2M \times N$ Klein bottle the partition function $Z_{2M,N}^{\text{Kln}}(K)$ is given by (36) in which the generating function $G^{\text{Kln}}(z, z, z)$ is given by (47). For a $(2M-1) \times N$ Klein bottle the partition function $Z_{2M-1,N}^{\text{Kln}}(K)$ is given by (50) in which the generating function $G^{\text{Kln}}(z, z_0, 1, z)$, $z_0 = \tanh(K/2)$, is computed using (51). All solutions yield the same Onsager bulk free energy (55).

We have also presented results of finite-size analyses of our solutions as well as those of the Ising model under cylindrical and toroidal boundary conditions at criticality. The analyses yield a central charge $c = 1/2$ in agreement with the conformal field prediction [4].

ACKNOWLEDGEMENT

Work has been supported in part by National Science Foundation Grant DMR-9980440.

- [7] B. M McCoy and T. T. Wu, *The Two-dimensional Ising Model*, (Harvard University Press, 1973).
- [8] P.W. Kasteleyn, Dimer statistics and phase transitions, *J. Math. Phys.* **4** (1963) 287-293.
- [9] P.W. Kasteleyn, The statistics of dimers on a lattice, *Physica* **27** (1961) 1209-1225.
- [10] W. T. Lu and F. Y. Wu, Partition function zeroes of a self-dual Ising model, *Physica A* **258** (1998) 157-170.
- [11] See, for example, K. Huang, *Statistical Mechanics*, (John Wiley and Son, New York, 1987), p. 387.
- [12] A. S. Ferdinand and M. E. Fisher, Bounded and inhomogeneous Ising models. I. Specific-heat anomaly of a finite lattice. *Phys. Rev.* **185** (1969) 832-846.
- [13] A. E. Ferdinand, Statistical mechanics of dimers on a quadratic lattice, *J. Math. Phys.* **8** (1967) 2332-2339.
- [14] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, (Academic Press, New York, 1994), 8.180.

-
- [1] W. T. Lu and F. Y. Wu, Dimer statistics on the Möbius strip and the Klein bottle. *Phys. Lett. A* **259** (1999) 108-114.
 - [2] W.-J. Tzeng, F. Y. Wu, Spanning trees on hypercubic lattices and nonorientable surfaces, *Appl. Math. Letters*, to appear, and preprint cond-mat/0001408.
 - [3] G. Tesler, Matchings in graphs and nonorientable surfaces, *J. Comb. Theory B* **78** (2000) 198-231.
 - [4] H. W. J. Blöte, J. L. Cardy, M. P. Nightingale, Conformal invariance, the central charge, and universal finite-size amplitudes at criticality, *Phys. Rev. Lett.* **56** (1986) 742-745.
 - [5] L. Onsager, Crystal statistics. I: A two-dimensional model with an order-disorder transition, *Phys. Rev.* **65** (1944) 117-149.
 - [6] B. Kaufman, Crystal statistics. II. Partition function evaluated by spinor analysis, *Phys. Rev.* **76** (1949) 1232-1243.